62000000201651
0523

Nothing works faster than the Paracetamol in Calpol

Formulation:

Each 5 mL of Strawberry-flavored alcohol free suspension contains 120 mg Paracetamol.

Use Paracetamol (Calpol) For:

- Relief of mild to moderate fever and pain
- Headache
- Migraine
- Muscle ache
- Sore throat
- Musculoskeletal pain
- Pain after dental procedure/tooth extraction
- Toothache
- Fever and pain associated with vaccination / immunization
- Ear ache / otalgia
- Respiratory tract infections including cold and flu

- Read full prescribing information before use. See inside panel.
- If you have any difficulties or do not understand the instructions ask your doctor or pharmacist.
- Patients who have been diagnosed with liver or kidney impairment must seek medical advice before taking this medication.
- If symptoms persist seek medical advice.
- Keep all medicines out of sight and reach of children.
- Store at temperatures not exceeding 30°C
- Keep out of direct light exposure.

Trade marks are owned by or licensed to the GSK group of companies.
©2021 GSK group of companies or its licensor

Strawberry Flavor

2-6 YEARS

gsk

Paracetamol
Calpol

120 mg/5 mL Suspension

Children

2-6 YEARS

Fast Fever Relief
Effective Relief of Pain

Strawberry Flavor   60 mL

Proof of Purchase

Paracetamol
Calpol

120 mg/5 mL Strawberry 60mL

How To Use Paracetamol (Calpol) 120 mg/5 mL

SHAKE WELL BEFORE USE   **READ TABLE FOR DOSAGE**

Check table for proper dosing

WEIGHT	DOSE (mL)
8 - 11 Kg (18 - 24 Lbs)	5
12 - 15 Kg (26 - 33 Lbs)	7.5
16 - 20 Kg (35 - 44 Lbs)	10
21 - 24 Kg (46 - 53 Lbs)	13

Based on recommended dosage of 10-15 mg/kg
- To be taken orally every 4-6 hours.
- Do not take more than 4 doses in any 24 hour period.
- Should not be used with other paracetamol-containing products.
- Maximum duration of continued use without medical advice: 3 days.

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph.
For first sign of adverse drug reaction, seek medical attention immediately.

📞 1800-14410884
📞 1800-89088275
✉ ph.customer-relations@gsk.com
🌐 www.gsk.com
DRHR-2132

---

gsk

Paracetamol
Calpol

120 mg/5 mL Suspension

Analgesic/Antipyretic

Manufactured by:
PT. Sterling Products Indonesia
Jl. Pulobuaran Raya,
Kav. III/DD 2,3,4 - Kawasan Industri Pulogadung,
Jakarta Timur, Indonesia

Imported by:
GlaxoSmithKline Consumer Healthcare Philippines, Inc.
23F The Finance Centre, 26th St. cor. 9th Ave.
Bonifacio Global City, Taguig City

4 800318 000190

62000000201651 0523

15°
48.5
106.5   234
47.5

BN
MFD
EXP

PHARMA CODE REF. IS:
8 2

36.93

Inside panel

gsk

Paracetamol
Calpol

Analgesic/Antipyretic

Paracetamol (Calpol) is available as:
- 120 mg/5 mL Suspension (60 mL, 100 mL)

What is in the medicine

- Each 5mL of Strawberry-flavored, alcohol-free **Paracetamol (Calpol)** Suspension with an odor characteristic of Strawberry contains 120 mg Paracetamol.

Research suggests that the maximum dose is more effective
- To be taken orally every 4-6 hours
- Minimum dosing interval: 4 hours
- Do not take more than 4 doses in any 24 hour period.
- Do not take for more than 3 days without asking your doctor.
- Do not exceed stated dose.
- Maximum Daily Dosage: 60 mg presented in divided doses of 10-15 mg/kg throughout the 24 hour period.
- Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary.

Contraindications, Precautions, Warnings (When should you not take this medicine)
- Contraindicated in patient with known hypersensitivity to paracetamol, or any other component of the formulation.
- Do not take more than the recommended dose as it may cause serious harm to your liver.
- If child has been diagnosed with liver or kidney impairment, seek medical advice before giving the medication.
- Check with your doctor before use if you:
  - have liver or kidney problems
  - are underweight or malnourished.

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.
  - have a severe infection as this may increase the risk of metabolic acidosis.
- Keep out of sight and reach of children.
- In patients with glutathione depleted states such as sepsis, the use of Paracetamol may increase the risk of metabolic acidosis.

Signs of metabolic acidosis include:
  - deep, rapid, difficult breathing
  - feeling sick (nausea), being sick (vomiting)
  - loss of appetite

When should you consult your doctor
- If you have any difficulties or do not understand the instructions ask your doctor or pharmacist.
- If symptoms persist seek medical advice.

What is the medicine used for

Used in the relief of mild to moderate fever and pain:
- Headache
- Migraine
- Muscle ache
- Sore throat
- Musculoskeletal pain
- Pain after dental procedure / tooth extraction
- Toothache
- Fever and pain associated with vaccination / immunization
- Ear ache / otalgia
- Respiratory tract infections including cold and flu

How To Use Paracetamol (Calpol)

Shake well before use and read table for dosage instructions.

WEIGHT	DOSE
Kg (Lbs)	120 mg/5 mL
8 - 11 (18 - 24)	5 mL
12 - 15 (26 - 33)	7.5 mL
16 - 20 (35 - 44)	10 mL
21 - 24 (46 - 53)	13 mL

Based on recommended dosage of 10-15 mg/kg

Interactions with Other Medications (What other medicine or food should be avoided when taking Calpol)
- The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.
- Should not be used with other Paracetamol containing products.

What should you do if you miss a dose
If fever/pain persists, give the missed dose and ensure 4 hours interval before giving the next dose.

Overdose (Care that should be taken when taking Calpol)
Paracetamol overdose may cause liver failure which may require liver transplant or lead to death. Acute pancreatitis has been observed, usually with hepatic dysfunction and liver toxicity.

What to do when you have taken more than the recommended dosage
- Immediate medical management is required in the event of overdose, even if symptoms of overdose are not present.
- Administration of N-acetylcysteine or methionine may be required.
- If overdose is confirmed or suspected, seek immediate advice from your Poison Centre and go to the nearest Emergency Medical Centre for management and expert treatment.

Undesirable Effects
Thrombocytopenia, anaphylaxis, cutaneous hypersensitivity reactions including skin rashes, angioedema and Stevens Johnson Syndrome, bronchospasm in patients sensitive to aspirin and other NSAIDs, hepatic dysfunction.

How to Store Calpol
Store at temperatures not exceeding 30°C

Reporting of side effects/adverse effects
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. For first sign of adverse drug reaction, seek medical attention immediately. You can also report side effects directly via consumer hotline.

📞 1800-89088275 / 1800-14410884
✉ ph.customer-relations@gsk.com
🌐 www.gsk.com

Bucknik et al: Comparison of Oral Versus Normal and High-Dose Rectal Acetaminophen in the Treatment of Febrile Children. Pediatrics 100, 553-556; 2002.

120 mg/5 mL Suspension
Amber PET bottle with CRSF cap (60 mL, 100 mL)
Registration No. DRHR-2132
Date of first Authorization: 31 Mar 2022
Date of Revision: May 2023
GD0v.7

---

gsk

Manufactured by: **PT. Sterling Products Indonesia**
Jl. Pulobuaran Raya, Kav. III/DD 2,3,4 - Kawasan Industri Pulogadung, Jakarta Timur
Imported by: **GlaxoSmithKline Consumer Healthcare Philippines, Inc.**
23F The Finance Centre, 26th St. cor. 9th Ave. Bonifacio Global City, Taguig City

62000000201651 0523

106.5   36.93
207
49
49
48.5   106.5   47.5
234